

REVUE SUISSE DE ZOOLOGIE

Tome 113 — Fascicule 1

	Pages
DANKITTIPAKUL, Pakawin, SONTICHAI, Saowapa & WANG, Xin Ping. Ten new species of coelotine spiders (Araneae, Amaurobiidae) from Thailand	3-21
LÖBL, Ivan. On the Philippine species of Cypariini and Scaphidiini (Coleoptera: Staphylinidae: Scaphidiinae)	23-49
TU, Lihong & LI, Shuqiang. A review of <i>Gongylidioides</i> spiders (Araneae: Linyphiidae: Erigoninae) from China	51-65
OSELLA, Giuseppe & ZUPPA, Anna Maria. <i>Otiorhynchus (Podonebistus) gasparoi</i> sp. n., un Curculionide anoftalmo della Grecia (Coleoptera, Curculionidae, Entiminae, Otiorhynchini)	67-75
SCHÄTTI, Beat. Northeast African racers of the <i>Platyceps rhodorachis</i> complex (Reptilia: Squamata: Colubrinae)	77-86
ALMIRÓN, Adriana E., AZPELIQUETA, M. de las Mercedes, CASCIOTTA, Jorge R. & LITZ, Thomas. A new species of <i>Hisonotus</i> (Siluriformes, Loricariidae, Otothyriini) from the República Oriental del Uruguay	87-94
MEREGALLI, Massimo, OSELLA, Giuseppe & ZUPPA, Anna Maria. The Raymondionymidae of the Curti collection, with description of <i>Raymondionymus curtii</i> sp. n. (Coleoptera, Curculionoidea)	95-113
EGUCHI, Katsuyuki. Six new species of <i>Pheidole</i> Westwood from North Vietnam (Hymenoptera, Formicidae)	115-131
INGRISCH, Sigfrid. New taxa and notes on some previously described species of scaly crickets from South East Asia (Orthoptera, Grylloidea, Mogoplistidae, Mogoplistinae)	133-227

REVUE SUISSE DE ZOOLOGIE
SWISS JOURNAL OF ZOOLOGY

tome 113, fascicule 1, mars 2006
Résumés

Ten new species of coelotine spiders (Araneae, Amaurobiidae) from Thailand

Pakawin DANKITTIPAKUL^{1, 2}, Saowapa SONTICHAI² & Xin Ping WANG³

¹ The University of Auckland, Private Bag 92019, Auckland, New Zealand.

E-mail: pdan021@ec.auckland.ac.nz

² Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand.

³ Research Associate at the Department of Entomology, California Academy of Sciences, San Francisco, CA 94118 and at the Florida State Collection of Arthropods, Gainesville, FL 32608, USA.

³ Coordinator of Statistical Research at the Brooks Center for Rehabilitation Studies, Health Science Center, University of Florida, P.O. Box 100185, Gainesville, FL 32610 and at the Rehabilitation Outcomes Research Center, Malcom Randall VA Medical Center, Gainesville, FL 32608, USA. E-mail: xwang@phhp.ufl.edu

Ten new species of coelotine spiders (Araneae, Amaurobiidae) from Thailand. - Ten new coelotine spiders of the genera *Coronilla* and *Draconarius* are described from Thailand: *C. lanna* sp. n. (♀), *D. australis* sp. n. (♂), *D. monticola* sp. n. (♀), *D. montis* sp. n. (♂), *D. phuhin* sp. n. (♂, ♀), *D. promontorius* sp. n. (♀), *D. schwendingeri* sp. n. (♂, ♀), *D. silva* sp. n. (♂), *D. silvicola* sp. n. (♀) and *D. tentus* sp. n. (♂). The genus *Coronilla* is recorded from Thailand for the first time. The species *D. australis* sp. n. marks the southernmost distribution of the subfamily Coelotinae.

Keywords: Coelotinae - *Coronilla* - *Draconarius* - new species - taxonomy - zoogeography - Thailand.

On the Philippine species of Cypariini and Scaphidiini (Coleoptera: Staphylinidae: Scaphidiinae)

Ivan LÖBL

Muséum d'histoire naturelle, Case postale 6434, CH-1211 Genève 6, Switzerland.

E-mail: ivan.lobl@mhn.ville-ge.ch

On the Philippine species of Cypariini and Scaphidiini (Coleoptera: Staphylinidae: Scaphidiinae). - The Philippine species of Cypariini and Scaphidiini are reviewed. The Cypariini are represented by a single species, *Cyparium punctatum* Pic, the Scaphidiini by 13 species of *Scaphidium*, with five species described as new: *S. crassipes*, *S. flavicorne*, *S. ilanum*, *S. kurozawai*, and *S. rufofemorale*. The previously described species of *Scaphidium* are redescribed, and a key to the species is given.

Keywords: Coleoptera - Staphylinidae - Scaphidiinae - Philippines - systematics.

A review of *Gongylidioides* spiders (Araneae: Linyphiidae: Erigoninae) from China

Lihong TU & Shuqiang LI

Institute of Zoology, Chinese Academy of Sciences, Beijing 100080, China.

Corresponding author: Shuqiang Li, e-mail: lisq@ioz.ac.cn

A review of *Gongylidioides* spiders (Araneae: Linyphiidae: Erigoninae) from China. - The present paper gives a review of the spider genus *Gongylidioides* occurring in China. A total of eight species are recognized, including three new species: *Gongylidioides acmodontus* sp. n., *Gongylidioides angustus* sp. n. and *Gongylidioides kouqianensis* sp. n. Descriptions of the new species and redescription of the known species, except for *G. onoi* and *G. ussuricus*, are provided.

Keywords: China - Linyphiidae - Erigoninae - *Gongylidioides* - new species - taxonomy.

***Otiorhynchus (Podonebistus) gasparoi* sp. n., un Curculionide anoftalmo della Grecia (Coleoptera, Curculionidae, Entiminae, Otiorhynchini)**

Giuseppe OSELLA & Anna Maria ZUPPA

Dipartimento di Scienze Ambientali, Università degli Studi dell'Aquila, Via Vetoio, I-67100 L'Aquila, Italia.

***Otiorhynchus (Podonebistus) gasparoi* n. sp., a blind weevil from Greece (Coleoptera, Curculionidae, Entiminae, Otiorhynchini).** - *Otiorhynchus (Podonebistus) gasparoi* sp. n. from Megalo Spiliò Monastiraki (Acarmania – Greece) is described. This new, blind species, collected in a cave, is very closely related to some eyeless *Otiorhynchus* from Epirus and Jonian islands (Greece): *O. (P.) doriae* (Solari & Solari, 1903), *O. (P.) imprevisus* Magnano, 1998 and *O. (P.) loebli* (Osella, 1974) (all included in the subgenus *Podonebistus* Reitter, 1912 of the genus *Otiorhynchus* Germar, 1824). *Otiorhynchus (Podonebistus) gasparoi* is easily distinguishable by pronotum subparallel-sided, metafemora with very large tooth, elytral punctures vanishing and interstriae with long half-lifted setae. The taxon, probably rizophagous on *Quercus coccifera* L. roots, was collected in the Megalo Spiliò cave jointly to other cave-dwelling arthropods, like the Pseudoscorpiones (an undescribed species of *Neobisium*), Isopoda, Orthoptera and as some Coleoptera Carabidae like *Duvalius (Eduvalius) ruffoanus* Casale, Giachino, Vailati & Vigna Taglianti, 1996 and *Laemostenus (Antisphodrus) giachinoi* Casale, 1997.

Keywords: Curculionidae - *Podonebistus* - new species - Greece.

Northeast African racers of the *Platyceps rhodorachis* complex (Reptilia: Squamata: Colubrinae)

Beat SCHÄTTI

Apartado postal 383, San Miguel de Allende, Gto. 37700, República Mexicana.

Northeast African racers of the *Platyceps rhodorachis* complex (Reptilia: Squamata: Colubrinae). - Racers from Eritrea and the Dahlak archipelago to central Somalia usually referred to *Platyceps rhodorachis subniger* (Boettger, 1893) are remarkable for the homogeneity of their scale features (e.g., preocular, anterior subocular, supralabials, ventrals, subcaudals, and dorsal scale rows). This taxon is morphologically closest to southern Arabian populations of Jan's cliff racer, *P. rhodorachis* auct.

Keywords: *Platyceps rhodorachis subniger* - *P. rhodorachis* auct. - *P. rhodorachis* group - morphology - distribution.

A new species of *Hisonotus* (Siluriformes, Loricariidae, Otothyridae) from the República Oriental del Uruguay

Adriana E. ALMIRÓN¹, M. de las Mercedes AZPELICUETA¹,
Jorge R. CASCIOTTA¹ & Thomas LITZ²

¹ División Zoología Vertebrados, Facultad de Ciencias Naturales y Museo, Paseo del Bosque s/n, 1900 La Plata, Argentina. E-mail: aalmiron@museo.fcnym.unlp.edu.ar

² Krumpfhalde 47, 88448 Attenweiler, Germany

A new species of *Hisonotus* (Siluriformes, Loricariidae, Otothyridae) from the República Oriental del Uruguay. - *Hisonotus charrua* sp. n. is described from the Río de la Plata and río Uruguay basins in the República Oriental del Uruguay. *Hisonotus charrua* sp. n. is distinguished by the following combination of characters: lateral plates 23-25, posterior margin of pectoral spine smooth, 18 to 24 plates bearing laterosensory canals, snout with an anterior odontode free area, and caudal peduncle depth 10.8-12.3% SL.

Keywords: Freshwaters - Siluriformes - loricarids - Hypoptopomatinae - *Hisonotus* - systematics.

The Raymondionymidae of the Curti collection, with description of *Raymondionymus curtii* sp. n. (Coleoptera, Curculionoidea)

Massimo MEREGALLI¹, Giuseppe OSELLA², Anna Maria ZUPPA²

¹ Dipartimento di Biologia Animale e dell'Uomo, Università di Torino, V. Accademia Albertina 13, I-10123 Torino, Italia.

E-mail: massimo.meregalli@unito.it

² Dipartimento di Scienze Ambientali, Università di L'Aquila, loc. Coppito, I-67100 L'Aquila, Italia. E-mail: osella@univaq.it

The Raymondionymidae of the Curti collection, with description of *Raymondionymus curtii* sp. n. (Coleoptera, Curculionoidea). - The Coleoptera Curculionoidea Raymondionymidae collected by Marc Curti and preserved in the Muséum d'histoire naturelle, Geneva, Switzerland were studied. A list of the 16 species and all specimens is given. Taxonomy, mutual relationships and distribution range of *R. ochsii*, *R. problematicus* and *R. orientalis* are discussed. *R. curtii* sp. n. (type locality: Italy, Piedmont, Valle Varaita, Castello, 44°37'N 07°03'E) is described. *R. sanfilippo* is a new record for the French fauna. Short remarks on the relationships among some species of the genus are also given.

Keywords: Coleoptera - Curculionoidea - Raymondionymidae - taxonomy - new species - Curti collection.

Six new species of *Pheidole* Westwood from North Vietnam (Hymenoptera, Formicidae)

Katsuyuki EGUCHI

The Kagoshima University Museum, Kagoshima University, Kagoshima 890-0065, Japan. E-mail: Katsuyuki.Eguchi@mc6.seikyoku.ne.jp

Six new species of *Pheidole* Westwood from North Vietnam (Hymenoptera, Formicidae). - Six new species of *Pheidole* Westwood are described from North Vietnam: *Pheidole colpigaleata* sp. nov., *Pheidole fortis* sp. nov., *Pheidole foveolata* sp. nov., *Pheidole laevicolor* sp. nov., *Pheidole magna* sp. nov. and *Pheidole vulgaris* sp. nov. *Pheidole fortis* was collected also from North Thailand, and *Pheidole vulgaris* is widespread through the Indo-Chinese subregion.

Keywords: Ant - *Pheidole* - new species - Vietnam.

New taxa and notes on some previously described species of scaly crickets from South East Asia (Orthoptera, Grylloidea, Mogoplistidae, Mogoplistinae)

Sigfrid INGRISCH

Zoologisches Forschungsinstitut & Museum Alexander Koenig (ZFMK),

Adenauerallee 160, D-53113 Bonn, Germany.

E-mail: sigfrid.ingrisc@planet-interkom.de

New taxa and notes on some previously described species of scaly crickets from South East Asia (Orthoptera, Grylloidea, Mogoplistidae, Mogoplistinae). - The Mogoplistinae of SE Asia are reviewed. A key to the genera in SE Asia is given. The genera are diagnosed. Two genera are described as new: *Apterornebius* gen. n. and *Terraplistes* gen. n. Keys to the species of the genera *Ornebius* Guérin-Méneville, 1844, *Ectatoderus* Guérin-Méneville, 1847, *Apterornebius* gen. n., *Gotvendia* Bolívar, 1927, *Cycloptiloides* Sjöstedt, 1909, and *Terraplistes* gen. n. are provided. The male phallic complex, male and female supra-anal plate, modification of the paraprocts, the ovipositor apical valves, and the maxillary palps are used as main diagnostic characters. 35 species are described as new: *Ornebius pullus* sp. n. (Brunei); *Ornebius cibodas* sp. n., *Ornebius samudra* sp. n., *Ornebius imitatus* sp. n., *Ornebius bogor* sp. n. (Java); *Ornebius citrus* sp. n. (Sabah); *Ornebius albipalpus* sp. n. (Singapore); *Ornebius dumoga* sp. n., *Ornebius consternus* sp. n. (Sulawesi); *Ornebius aureus* sp. n., *Ornebius serratus* sp. n., *Ornebius angustus* sp. n., *Ornebius tuberculatus* sp. n., *Ornebius peniculatus* sp. n., *Ornebius brevipalpus* sp. n. (Thailand); *Apterornebius kinabalu* sp. n. (Sabah); *Apterornebius chong* sp. n. (Thailand);

Ectatoderus samui sp. n., *Ectatoderus argentatus* sp. n. (Thailand); *Gotwendia erawan* sp. n. (Thailand); *Micrornebius lineatus* sp. n. (Sabah); *Micrornebius cylindricus* sp. n. (Singapore); *Micrornebius maninjau* sp. n. (Sumatra); *Micrornebius insularis* sp. n., *Micrornebius laem* sp. n., *Micrornebius inopinatus* sp. n. (Thailand); *Cycloptiloides timah* sp. n. (Singapore); *Cycloptiloides pui* sp. n., *Cycloptiloides pakchong* sp. n., *Cycloptiloides lobicauda* sp. n. (Thailand); *Terraplistes chantri* sp. n., *Terraplistes kradung* sp. n., *Terraplistes erawan* sp. n., *Terraplistes brevicauda* sp. n., and *Terraplistes excisa* sp. n. (Thailand). Four new combinations are proposed: *Micrornebius sandrasagarai* (Fernando, 1957) comb. n. from *Ectatoderus*, *Micrornebius brevipalpis* (Chopard, 1930) comb. n. from *Ornebius*, *Micrornebius incertus* (Ingrisch, 1998) comb. n. from *Derectaotus*, *Terraplistes niger* (Ingrisch, 1987) comb. n. from *Cycloptiloides*; [*Ectatoderus pallidigeniculatus* Brunner, 1893 probably also belongs to *Micrornebius*]. Types of the new species are in the museums in Bonn, Geneva or London. Notes on some previously described species are given. For five species (*Ornebius aureus*, *O. samudra*, *Ectatoderus argentatus*, *Micrornebius inopinatus*, and *M. maninjau*) stridulation is also described.

Keywords: Grylloidea - Mogoplistinae - South East Asia - diagnostic characters - new taxa - keys to genera and species.